

inert carriers and other components to facilitate handling and application for particular target pests. The formulation and application procedures are all well known in the art and are used with commercial strains of *B. thuringiensis* (HD-1) active against Lepidoptera, e.g., caterpillars.

5

4.3 RECOMBINANT HOST CELLS FOR EXPRESSING THE *CRYIC** GENES

The nucleotide sequences of the subject invention can be introduced into a wide variety of microbial hosts. Expression of the toxin gene results, directly or indirectly, in the intracellular production and maintenance of the pesticide. With suitable hosts, e.g., *Pseudomonas*, the microbes can be applied to the sites of lepidopteran insects where they will proliferate and be ingested by the insects. The result is a control of the unwanted insects. Alternatively, the microbe hosting the toxin gene can be treated under conditions that prolong the activity of the toxin produced in the cell. The treated cell then can be applied to the environment of target pest(s). The resulting product retains the toxicity of the *B. thuringiensis* toxin.

10

15

Suitable host cells, where the pesticide-containing cells will be treated to prolong the activity of the toxin in the cell when the then treated cell is applied to the environment of target pest(s), may include either prokaryotes or eukaryotes, normally being limited to those cells which do not produce substances toxic to higher organisms, such as mammals. However, organisms which produce substances toxic to higher organisms could be used, where the toxin is unstable or the level of application sufficiently low as to avoid any possibility or toxicity to a mammalian host. As hosts, of particular interest will be the prokaryotes and the lower eukaryotes, such as fungi. Illustrative prokaryotes, both Gram-negative and Gram-positive, include *Enterobacteriaceae*, such as *Escherichia*, *Erwinia*, *Shigella*, *Salmonella*, and *Proteus*; *Bacillaceae*; *Rhizobiceae*, such as *Rhizobium*; *Spirillaceae*, such as photobacterium, *Zymomonas*, *Serratia*, *Aeromonas*, *Vibrio*, *Desulfovibrio*, *Spirillum*; *Lactobacillaceae*; *Pseudomonadaceae*, such as *Pseudomonas* and *Acetobacter*; *Azotobacteraceae*, *Actinomycetales*, and *Nitrobacteraceae*. Among eukaryotes are fungi, such as *Phycomycetes* and *Ascomycetes*, which includes yeast, such

20

25

as *Saccharomyces* and *Schizosaccharomyces*; and *Basidiomycetes* yeast, such as *Rhodotorula*, *Aureobasidium*, *Sporobolomyces*, and the like.

Characteristics of particular interest in selecting a host cell for purposes of production include ease of introducing the *B. thuringiensis* gene into the host, availability of expression systems, efficiency of expression, stability of the pesticide in the host, and the presence of auxiliary genetic capabilities. Characteristics of interest for use as a pesticide microcapsule include protective qualities for the pesticide, such as thick cell walls, pigmentation, and intracellular packaging or formation of inclusion bodies; leaf affinity; lack of mammalian toxicity; attractiveness to pests for ingestion; ease of killing and fixing without damage to the toxin; and the like. Other considerations include ease of formulation and handling, economics, storage stability, and the like.

Host organisms of particular interest include yeast, such as *Rhodotorula sp.*, *Aureobasidium sp.*, *Saccharomyces sp.*, and *Sporobolomyces sp.*; phylloplane organisms such as *Pseudomonas sp.*, *Erwinia sp.* and *Flavobacterium sp.*; or such other organisms as *Escherichia*, *Lactobacillus sp.*, *Bacillus sp.*, *Streptomyces sp.*, and the like. Specific organisms include *Pseudomonas aeruginosa*, *Pseudomonas fluorescens*, *Saccharomyces cerevisiae*, *Bacillus thuringiensis*, *Escherichia coli*, *Bacillus subtilis*, *Bacillus megaterium*, *Bacillus cereus*, *Streptomyces lividans* and the like.

Treatment of the microbial cell, e.g., a microbe containing the *B. thuringiensis* toxin gene, can be by chemical or physical means, or by a combination of chemical and/or physical means, so long as the technique does not deleteriously affect the properties of the toxin, nor diminish the cellular capability in protecting the toxin. Examples of chemical reagents are halogenating agents, particularly halogens of atomic no. 17-80. More particularly, iodine can be used under mild conditions and for sufficient time to achieve the desired results. Other suitable techniques include treatment with aldehydes, such as formaldehyde and glutaraldehyde; anti-infectives, such as zephiran chloride and cetylpyridinium chloride; alcohols, such as isopropyl and ethanol; various histologic fixatives, such as Lugol's iodine, Bouin's fixative, and Helly's fixatives, (see e.g., Humason, 1967); or a combination of physical (heat) and chemical agents that preserve and prolong the activity of the toxin produced in the cell when the cell is

administered to the host animal. Examples of physical means are short wavelength radiation such as γ -radiation and X-radiation, freezing, UV irradiation, lyophilization, and the like. The cells employed will usually be intact and be substantially in the proliferative form when treated, rather than in a spore form, although in some instances spores may be employed.

Where the *B. thuringiensis* toxin gene is introduced via a suitable vector into a microbial host, and said host is applied to the environment in a living state, it is essential that certain host microbes be used. Microorganism hosts are selected which are known to occupy the "phytosphere" (phylloplane, phyllosphere, rhizosphere, and/or rhizoplane) of one or more crops of interest. These microorganisms are selected so as to be capable of successfully competing in the particular environment (crop and other insect habitats) with the wild-type microorganisms, provide for stable maintenance and expression of the gene expressing the polypeptide pesticide, and, desirably, provide for improved protection of the pesticide from environmental degradation and inactivation.

A large number of microorganisms are known to inhabit the phylloplane (the surface of the plant leaves) and/or the rhizosphere (the soil surrounding plant roots) of a wide variety of important crops. These microorganisms include bacteria, algae, and fungi. Of particular interest are microorganisms, such as bacteria, e.g., genera *Bacillus*, *Pseudomonas*, *Erwinia*, *Serratia*, *Klebsiella*, *Zanthomonas*, *Streptomyces*, *Rhizobium*, *Rhodopseudomonas*, *Methylophilus*, *Agrobacterium*, *Acetobacter*, *Lactobacillus*, *Arthrobacter*, *Azotobacter*, *Leuconostoc*, and *Alcaligenes*; fungi, particularly yeast, e.g., genera *Saccharomyces*, *Cryptococcus*, *Kluyveromyces*, *Sporobolomyces*, *Rhodotorula*, and *Aureobasidium*. Of particular interest are such phytosphere bacterial species as *Pseudomonas syringae*, *Pseudomonas fluorescens*, *Serratia marcescens*, *Acetobacter xylinum*, *Agrobacterium tumefaciens*, *Rhodobacter sphaeroides*, *Xanthomonas campestris*, *Rhizobium melioli*, *Alcaligenes eutrophus*, and *Azotobacter vinlandii*; and phytosphere yeast species such as *Rhodotorula rubra*, *R. glutinis*, *R. marina*, *R. aurantiaca*, *Cryptococcus albidus*, *C. diffluens*, *C. laurentii*, *Saccharomyces rosei*, *S. pretoriensis*, *S. cerevisiae*, *Sporobolomyces roseus*, *S. odorus*, *Kluyveromyces veronae*, and *Aureobasidium pollulans*.